


# LUX, and the Combating of the Lamppost Effect



*Matthew Szydagis, APS DPF Meeting at Fermilab, July 31, 2017*




# A Big Hole in Our Knowledge

Bill Watterson



## What is this dark matter?

<http://cdn.phys.org/newman/gfx/news/pires/2015/the-darksideo.png>




- Neutrinos 0.1%
  - Photons 0.01%
  - Black holes 0.005%
- WIMPs? (Weakly Interacting Massive Particles) Not this**

Above credit: X-ray: NASA/CXC/CfA/M. Markevitch et al.; Lensing Map: NASA/STScI; ESO WFI; Magellan/U. Arizona / D. Clowe et al.; Optical image: NASA / STScI; Magellan / U. Arizona / D. Clowe et al.; Right: NASA/ESA / M. Bradac et al.


# Large Underground Xenon

- \* 2-phase xenon detector deployed (was recently decommissioned) underground in SD with 122 photo-multiplier tubes
- \* Why element Xe?
  - \* Dense (good self-shielding)
  - \* Gets excited and scintillates, and can get ionized easily
- \* Why deep underground?
  - \* Cosmic rays -> bad
- \* Why PMTs not low BG Si?
  - \* PMT: 1-photon, large area
  - \* SiPM unavailable in past



photos by Jack Genovesi, UAlbany



# How It Functions



S2/S1 ratio gives particle ID,  
and S2-S1 drift time gives depth

- \* Two scintillation pulses, S1 and S2 (vacuum-UV)
- \* S1 in liquid + S2 in gas
- \* S1 O(10-100) ns-wide exponential, S2 O(1 microsec.) Gaussian
- \* S1 is direct photon counting, but S2 secondary photons from ionization e<sup>-</sup>'s
- \* Why 2 (forms of light)?
  - \* Better position and energy reconstruction
  - \* Particle identification
  - \* Reuse the same PMTs


# Position Reconstruction




- \* Even a single drift electron from an ionization is visible using the S2!
- \* X-Y position is reconstructed at 2-20 mm accuracy using top PMTs
  - \* Depends on S2 size, and on radius
  - \* Detector was 50x50cm dia x depth
- \* Possible to reconstruct positions of neutron elastic scatters from D-D gun, and isotropic internal sources
  - \* Neutrons like dark matter WIMP signal in theory, therefore used for the calibrations
  - \*  $\text{CH}_3\text{T}$ ,  $^{83\text{m}}\text{Kr}$ ,  $^{127}\text{Xe}$  calibrate the BGs
- \* Many more technical publications forthcoming (and physics results!)

# Calibration: NR With D-D Gun

- \* Lowest absolute calibration of light yield (180 V/cm)
  - \* 1.1 keVnr
  - \* Previous 3 keVnr (from Plante et al., 2011) 0 field
- \* Lowest absolute, direct calibration of charge yield (180 V/cm)
  - \* 0.6 keVnr
  - \* Previous was actually only 4 keVnr! (from Manzur et al., 2010)
- \* Air-filled conduit in water shield is neutron guide




# Calibrations: ER, Old and New




tritium (i.e.,  
tritiated  
methane)




$^{83m}\text{Kr}$




low energy x-rays  
and electron  
capture or internal  
conversion events




# Efficiencies (Analysis, NR)



- 50% efficient @3-4 keVnr (depending on run)
- Below 1.1 keV ( $L_y$  un-calibrated) set to 0






# Efficiencies (Trigger, ER & NR)



The trigger thresholds are of course well below the analysis thresholds



# E-Field Modeling

## Wall Position: 2 Examples




# SI WIMP Search Final Results

(in S2/S1 space; limit on next slide)





Phys. Rev. Lett. 116, 161301 (2016), re-analysis of  
Phys. Rev. Lett. 112, 091303 (2014)

Phys. Rev. Lett. 118, 021303 (2017)



# Spin-dependent Limits

Phys. Rev. Lett. 118, 251302 (2017)




- Left plot is neutron coupling vs. mass, while right is proton interaction strength vs. neutron, at a fixed example mass near the strictest point in the limit curve (50 GeV)
- Xe is even Z, but some isotopes are odd-N, allowing for SD interactions to be probed, especially WIMP-neutron: LHC dark matter limits exceeded at high mass


# No WIMPs, So Trying Axions

*First Run Only*



Solar Axions





Galactic ALPs



Phys. Rev. Lett. 118, 261301 (2017)



# Push to High-E WIMPs (for *EFT*)



# Conclusions

- \* The LUX spin-independent WIMP limit led the field for 3 years (2013-2016). Only now are the larger XeTPCs catching up (XENON1T, Panda-X) ☺ arXiv:1705.06655, arXiv:1707.07921
- \* LUX ultimately delivered \*4\* times better sensitivity in 427 live-days than projected 300 live-day sensitivity for design in the original LUX proposal
  - \* This is nearly unheard of, especially in direct WIMP dark matter searches!
- \* Spin-dependent limit still best in world for neutrons
- \* Strictest constraints on axions and axion-like particles in terms of coupling to electrons
- \* Pushing on combining PSD from S1 with S2/S1 discrimination, to use effectively for first time in LXeTPC (Effective Field Theory analysis soon)
- \* LUX yields, efficiencies, and fields well calibrated, simulated, and understood, for all runs
- \* LUX is not done yet: lot more papers to come out of data!
  - \* There is a great deal more science yet to come. Be on the look out



Hopefully we are looking for dark matter in ALL the right places !

# Thank You! Questions??




Honoré Daumier, "Mr. Babinet, warned by his concierge of the arrival of the comet", illustration for *Le Charivari*, 22 September 1858

# SD Proton, and Different Example Mass for $a_p$ v. $a_n$



# Nod to XENON1T's First



# Primordial BHs as DM???




FIG. 3: Constraints on  $f(M)$  for a variety of evaporation (magenta), dynamical (red), lensing (cyan), large-scale structure (green) and accretion (orange) effects associated with PBHs. The effects are extragalactic  $\gamma$ -rays from evaporation (EG) [11], femtolensing of  $\gamma$ -ray bursts (F) [187], white-dwarf explosions (WD) [188], neutron-star capture (NS) [36], Kepler microlensing of stars (K) [189], MACHO/EROS/OGLE microlensing of stars (ML) [27, 190] and quasar microlensing (broken line) (ML) [191], survival of a star cluster in Eridanus II (E) [192], wide-binary disruption (WB) [37], dynamical friction on halo objects (DF) [33], millilensing of quasars (mLQ) [32], generation of large-scale structure through Poisson fluctuations (LSS) [14], and accretion effects (WMAP, FIRAS) [15]. Only the strongest constraint is usually included in each mass range, but the accretion limits are shown with broken lines since they are highly model-dependent. Where a constraint depends on some extra parameter which is not well-known, we use a typical value. Most constraints cut off at high  $M$  due to the incredulity limit. See the original references for more accurate forms of these constraints.

# Pathologies

