



# A Strategy for LATTICE FIELD THEORY on curved RIEMANN MANIFOLDS




REGGE

+



FINITE ELEMENT (FEM)

+




QUANTUM CENTER TERM = QFE

Rich Brower, Boston University: Paris June 12, 2018  
with G. Fleming, A. Gasbarro, T. Raben, C-I Tan, E. Weinberg

<https://arxiv.org/abs/1610.08587> Dirac Fermions on Simplicial Manifold  
<https://arxiv.org/abs/1803.08512> Phi 4th on Riemannian Manifold

# BREAKING OUT OF FLAT LAND LATTICE FIELD THEORY



# QUANTUM FINITE ELEMENT (QFE) CONSTRUCTION

## 1. “Regular” Simplicial Complex (aka triangulation)

- Curvature  $R = 0, R > 0, R < 0$

## 2. Finite Elements/Regge for Free Fields


- Spin:  $J = 0$  (scalar),  $1/2$  (Dirac),  $1$  (Gauge)

## 3. UV problem and Quantum Corrections

- $c = 1/2$  CFT (aka  $\phi^4$  Ising) on  $S^2$

# REGULAR TRIANGULATIONS!

Triangle case



Preserves Discrete  
Subgroup of Isometries

$$\frac{1}{p} + \frac{1}{q} > 1/2$$

de Sitter  $S^2$

vertex  $q = 3, 4, 5$

$$\frac{1}{p} + \frac{1}{q} = 1/2$$

flat  $T^2$

vertex  $q = 6$


$$\frac{1}{p} + \frac{1}{q} < 1/2$$

Hyperbolic  $AdS^2$



vertex  $q = 7, 8, 9, \dots$

# *Start with maximum regular Tessellation*

$$s = 1$$



$$s = 8$$




$I = 0$  (A),  $1$  (T1) ,  $2$  (H) are irreducible 120 Icosahedral subgroup of  $O(3)$


$1/p + 1/q > 1/2$  for regular positive curvature tessellation

# Sphere Constant Positive Curvature vs Cylinders

$$\mathbb{S}^d \rightarrow \mathbb{R} \times \mathbb{S}^d$$



Aristotle's 2% Error!





Fast Code Domains of  
Regular 3D Grids on Refinement

$$(2\pi - 5\text{ArcCos}[1/3]) / (2\pi) = 0.0204336$$

The full **symmetry group** of the 600-cell is the **Weyl group** of  $H_4$ . This is a **group** of order 14400. It consists of 7200 **rotations** and 7200 rotation-reflections. The rotations form an **invariant subgroup** of the full symmetry group.

# Hyperbolic (e.g. Poincare Disk) and Global AdS

$$\mathbb{H}^d \rightarrow \text{AdS}^d$$



$$1/p + 1/q < 1/2$$

Triangle Group Tessellation:  
Preserve Finite subgroup of the Modular Group


These Hyperbolic Tessellations are "Tensor Networks":  
What can we do with them as lattice Field Theories?

# PART I: FREE THEORIES



## REGGE: Piecewise linear metric


$$(\mathcal{M}, g_{\mu\nu}(x)) \leftrightarrow (\mathcal{M}_\sigma, g_\sigma = \{l_{ij}\})$$



Simplicial Complex/Delaunay Dual Complex +  
Regge flat metric on each Simplex



## FEM: Piecewise linear fields

$$\phi(x) \leftrightarrow \phi = \sum_i \phi_i W_i(\xi)$$



Actually fancier methods: Discrete Exterior Calculus  
(scalar), Spin connection (Fermion), Wilson links  
(gauge), etc.

# REGGE CALCULUS FEM FORMULATION



LINEAR FEM/ REGGE CALCULUS \*

$$FEM : A_d \frac{(\phi_2 - \phi_3)^2}{l_1^2} \quad *d * d\phi_i$$

Delaunay Link Area:  $A_d = h_1 l_1$

DISCRETE EXTERIOR CALCULUS or CHRIST FRIEBERG & LEE

# SIMPLICIAL ALGEBRA

Building a Simplicial Complex  $\mathcal{S} \quad \sigma_0 \rightarrow \sigma_1 \rightarrow \sigma_2 \rightarrow \cdots \rightarrow \sigma_D$

Circumcenter Dual Lattice  $\mathcal{S}^* \quad \sigma_0^* \leftarrow \sigma_1^* \leftarrow \sigma_2^* \leftarrow \cdots \leftarrow \sigma_D^*$

boundary operator: 
$$\partial\sigma_n(i_0i_1 \cdots i_n) = \sum_{k=0}^n (-1)^k \sigma_{n-1}(i_0i_1 \cdots \widehat{i}_k \cdots i_n)$$

Orthogonality and proper hybrid tiling : 
$$V_{nn^*} = \langle \sigma_n | \sigma_n^* \rangle = \int \sigma_n \wedge \sigma_n^* = \frac{n!(D-n)!}{D!} |\sigma_n| |\sigma_n^*|$$

Add matter  $n$  form  $\omega_n$  on  $\sigma_n$

Stokes Theorem for Exterior Derivative

$$\int_{\sigma_n} d\omega(y) = \int_{\partial\sigma_n} \omega(y) \quad \text{or} \quad \langle \sigma_n | d\omega \rangle = \langle \partial\sigma_n | \omega \rangle$$


$$\delta = *d* \quad \implies \quad \delta^2 = d^2 = 0$$

Beltrami-Laplace  $(\delta + d)^2 \phi = *d * d\phi$       Kahler-Dirac  $(\delta + d)\psi = 0$

# FEM FIXES SPECTRAL DEFECTS OF LAPLACIAN ON SPHERE


For  $s = 8$  first  $(l+1)^2 = 64$  eigenvalues

BEFORE FEM






$l, m$

AFTER FEM



$l, m$

# SPECTRUM OF FEM LAPLACIAN ON A SPHERE



Fit

$$l + 1.00012 l^2$$

$$- 13.428110^{-6} l^3 - 5.5724410^{-6} l^4$$



# Dirac ON SIMPLICIAL MANIFOLD

$$S = \frac{1}{2} \int d^D x \sqrt{g} \bar{\psi} [\mathbf{e}^\mu (\partial_\mu - \frac{i}{4} \omega_\mu(x)) + m] \psi(x)$$

$$\mathbf{e}^\mu(x) \equiv e_a^\mu(x) \gamma^a \quad \text{Verbein \& Spin connection}^*$$

$$\omega_\mu(x) \equiv \omega_\mu^{ab}(x) \sigma_{ab} \quad , \quad \sigma_{ab} = i[\gamma_a, \gamma_b]/2$$


- (1) New spin structure “knows” about intrinsic geometry
- (2) Need to avoid simplex curvature singularities at sites.
- (3) Spinors rotations (Lorentz group) is double of  $O(D)$ .

$$e^{i(\theta/2)\sigma_3/2} \rightarrow -1 \quad \text{as} \quad \theta \rightarrow 2\pi$$

\* Must satisfy the tetrad postulate!

$$\omega_\mu^{ab} = \frac{1}{2} e^{\nu[a} (e_{\nu,\mu}^{b]} - e_{\mu,\nu}^{b]} + e^{b]\sigma} e_{\mu}^c e_{\nu c, \sigma}$$

$$S_{naive} = \frac{1}{2} \sum_{\langle i,j \rangle} \frac{V_{ij}}{l_{ij}} [\bar{\psi}_i \vec{e}^{(i)j} \cdot \vec{\gamma} \Omega_{ij} \psi_j - \bar{\psi}_j \Omega_{ji} \vec{e}^{(i)j} \cdot \vec{\gamma} \psi_i] + \frac{1}{2} m V_i \bar{\psi}_i \psi_i$$



**DO NOT USE PIECEWISE LINEAR REGGE CALCULUS MANIFOLD!**

Simplicial Tetrad Hypothesis

$$e_a^{(i)j} \gamma^a \Omega_{ij} + \Omega_{ij} e_a^{(j)i} \gamma^a = 0$$

Gauge Invariance under Spin(D) transformations

$$\psi_i \rightarrow \Lambda_i \psi \quad , \quad \bar{\psi}_j \rightarrow \bar{\psi}_j \Lambda_j^\dagger \quad , \quad \mathbf{e}^{(i)j} \rightarrow \Lambda_i \mathbf{e}^{(i)j} \Lambda_i^\dagger \quad , \quad \Omega_{ij} \rightarrow \Lambda_i \Omega_{ij} \Lambda_j^\dagger$$

# QFE SUMMARY OF SIMPLICIAL FIELD

$$\mathbf{J} = 0 \quad S_{scalar} = \frac{1}{2} \sum_{\langle i,j \rangle} \frac{V_{ij}}{l_{ij}^2} (\phi_i - \phi_j)^2 ,$$

$$\mathbf{J} = 1/2 \quad S_{Wilson} = \frac{1}{2} \sum_{\langle i,j \rangle} \frac{V_{ij}}{l_{ij}} (\bar{\psi}_i \hat{e}_a^{j(i)} \gamma^a \Omega_{ij} \psi_j - \bar{\psi}_j \Omega_{ji} \hat{e}_a^{i(j)} \gamma^a \psi_i)$$

$$\mathbf{J} = 1 \quad S_{gauge} = \frac{1}{2g^2 N_c} \sum_{\Delta_{ijk}} \frac{V_{ijk}}{A_{ijk}^2} Tr[2 - U_{\Delta_{ijk}} - U_{\Delta_{ijk}}^\dagger]$$



$$V_{ij} = |\sigma_2(ij) \wedge \sigma_2^*(ij)|$$

$$U_{\Delta_{ijk}} = U_{ij} U_{jk} U_{ki} \quad A_{ijk} = |\sigma_3(ijk)| \quad V_{ijk} = |\sigma_3(ijk) \wedge \sigma_3^*(ijk)|$$

$$\mathbf{U(1) QED} \quad 2 - U_{\Delta_{123}} - U_{\Delta_{123}}^\dagger = 2[1 - \cos(\theta_{12} + \theta_{23} + \theta_{31})] \\ \simeq (\theta_{12} + \theta_{23} + \theta_{31})^2$$



# 2D DIRAC SPECTRA ON SPHERE



$$E = j + 1/2$$

Exact is integer spacing for  $j = 1/2, 3/2, 5/2 \dots$  Exact degeneracy  $2j + 1$ : No zero mode in chiral limit!.

<https://arxiv.org/abs/1610.08587>




**Lattice Dirac Fermions on a Simplicial Riemannian Manifold**

Richard C. Brower, George T. Fleming, Andrew D. Gasbarro, Timothy G. Raben, Chung-I Tan, Evan S. Weinberg

# FREE MAJORANA FERMIONS ON S<sup>2</sup>


$$\langle \psi(z_1) \bar{\psi}(z_1) \bar{\psi}(z_1) \psi(z_2) \rangle = \left[ \frac{1}{\partial} \right]_{z_1, z_2} \left[ \frac{1}{\bar{\partial}} \right]_{z_1, z_2} = \frac{1}{4\pi^2} \frac{1}{|z_1 - z_2|^2}$$

$$\frac{\langle \sigma(0) \epsilon(z_2) \epsilon(z_3) \sigma(\infty) \rangle}{\langle \epsilon(z_2) \epsilon(z_3) \rangle} = \frac{1}{4} \left| \sqrt{z_1/z_2} + \sqrt{z_2/z_1} \right|^2 = \frac{1}{4} (r + 1/r + 2 \cos \theta)$$



# PARTII: INTERACTION THEORIES


TEST CFT: PHI 4TH AT WILSON-FISHER FIXED POINT IN 2D & 3D.



$$\mathcal{L}(x) = -\frac{1}{2}(\nabla\phi)^2 + \lambda(\phi^2 - \mu^2/2\lambda)^2$$

# TEST 2D ISING/PHI 4<sup>TH</sup> ON THE RIEMANN SPHERE

Stereographic project of Complex Plane:



$$\xi = \tan(\theta/2)e^{-i\phi} = \frac{x + iy}{1 + z}$$

$$|\xi| = \sqrt{\xi_1^2 + \xi_2^2} \quad \xi = \xi_1 + i\xi_2$$

$$\vec{r} = (x, y, z) \quad \vec{r} \cdot \vec{r} = 1$$

$$|\vec{r}_1 - \vec{r}_2| = 2 - 2 \cos(\theta_{12})$$


Conformally Invariant  
Cross Ratios are "Preserved"

$$\frac{|\xi_1 - \xi_2||\xi_3 - \xi_4|}{|\xi_1 - \xi_3||\xi_1 - \xi_4|} = \frac{|\vec{r}_1 - \vec{r}_2||\vec{r}_1 - \vec{r}_2|}{|\vec{r}_1 - \vec{r}_3||\vec{r}_1 - \vec{r}_4|}$$

# BINDER CUMULANT NEVER CONVERGES

$$U_4 = \frac{3}{2} \left[ 1 - \frac{\langle M^4 \rangle}{3 \langle M^2 \rangle^2} \right]$$

0.85102




Very fast cluster algorithm:

Brower, Tamayo 'Embedded Dynamics for phi 4<sup>th</sup> Theory" PRL 1989.


Wolff single cluster + plus Improved Estimators etc

# Restoring Isometries for ON A SIMPLICIAL COMPLEX

How much help do you need from FEM ?



Hypercubic Lattice




Simplicial Sphere

UV


IR

Is Renormalized perturbation theory at the UV fixed point enough to uniquely/correctly define the IR Quantum Field Theory?


# UV DIVERGENCE BREAKS ROTATIONS



$$\delta m^2 = \lambda \langle \phi(x) \phi(x) \rangle \rightarrow \frac{1}{K_{xx}}$$



one configuration



average of config.



# One LOOP Counter Term

---


$$\Delta m_i^2 = 6\lambda [K^{-1}]_{ii} \simeq \frac{\sqrt{3}}{8\pi} \lambda \log(1/m_0^2 a_i^2) = \frac{\sqrt{3}}{8\pi} \lambda \log(N_s) + \frac{\sqrt{3}}{8\pi} \lambda \log(a^2/a_i^2)$$

Exact Continuum  
Divergence

Local Cut-off  
Scheme Dependence

$$\delta\mu_i^2 = -6\lambda \left( [K^{-1}]_{ii} - \frac{1}{N_s} \sum_{j=1}^{N_s} [K^{-1}]_{jj} \right)$$

# MODEL OF COUNTER TERM



# RG Proof Of UNIVERSAL UV Logs

$$G_{xx}(m) \simeq c_x \log(1/m^2 a_x^2) + O(a^2 m^2)$$

$$\implies \gamma(m_0^2) = -m_0 \frac{\partial}{\partial m_0} G_{xx}(m_0^2) \simeq 2c_x + O(m_0^2)$$

$$m_0 = am \rightarrow 0$$

## FEM Spectral Fidelity

$$G_{xy}(m^2) = \sum_n \frac{\phi_n^*(x)\phi_n(x)}{E_n^{(0)} + m^2}$$

IR: Region.      UV





$$\simeq \frac{\sqrt{3}}{8\pi} \sum_{l=0}^{L_0} \frac{(2l+1)P_l(r_x \cdot r_y)}{l(l+1) + \mu_0^2} + \sum_{n=(L_0+1)^2}^N \frac{\phi_n^*(x)\phi_n(y)}{E_n^{(0)} + m^2}$$

## Insensitive to UV defects



$$\gamma(m_0^2) = -m_0 \frac{\partial}{\partial m_0} G_{xx}(m_0^2)$$

$$\simeq \frac{\sqrt{3}}{8\pi} \int_0^{\Lambda_0^2} dE^{(0)} \frac{m_0^2}{(E^{(0)} + m_0^2)^2} = \frac{\sqrt{3}}{8\pi} \frac{1}{1 + m_0^2/\Lambda_0^2}$$


# One Loop Counter Term vs Two Loop Convergence



# Counter term in 3D



$$\delta\mu_{CT}^2(x, y) \sim \lambda_0 c_x \delta_{xy} + \lambda_0^2 e^{-6|x-y|/a}$$



# EXACT C = 1/2 CFT ON 2D SPHERE

## Exact Two point function

$$\langle \phi(x_1)\phi(x_2) \rangle = \frac{1}{|x_1 - x_2|^{2\Delta}} \rightarrow \frac{1}{|1 - \cos\theta_{12}|^\Delta}$$

$$\Delta = \eta/2 = 1/8 \qquad x^2 + y^2 + z^2 = 1$$

4 pt function  $(x_1, x_2, x_3, x_4) = (0, z, 1, \infty)$

$$g(0, z, 1, \infty) = \frac{1}{2|z|^{1/4}|1-z|^{1/4}} [ |1 + \sqrt{1-z}| + |1 - \sqrt{1-z}| ]$$

Critical Binder Cumulant  $U_B^* = 1 - \frac{\langle M^4 \rangle}{3\langle M^2 \rangle \langle M^2 \rangle} = 0.567336$

Dual to Free Fermion

# Now Binder Cumulant Converges

$$U_{2n}(\mu^2, \lambda, s) = U_{2n,cr} + a_{2n}(\lambda)[\mu^2 - \mu_{cr}^2]s^{1/\nu} + b_{2n}(\lambda)s^{-\omega}$$

**FIT**  $U_{4,cr} = 0.85020(58)(90)$

**THEORY**  $U_4^* = 0.8510207(63)$


**FIT**  $U_{6,cr} = 0.77193(37)(90)$

**THEORY**  $U_6^* = 0.773144(21)$

$$U_4 = \frac{3}{2} \left[ 1 - \frac{\langle M^4 \rangle}{3 \langle M^2 \rangle \langle M^2 \rangle} \right]$$


$$\mu_{cr}^2 = 1.82240070(34)$$

$$dof = 1701 \quad , \quad \chi^2/dof = 1.026$$




Simultaneous fit for s up 800:  
E.G. 6,400,002 Sites on Sphere

$$\lambda_0 = 1.0, \mu_0^2 = 1.823405$$



$$\lambda_0 = 1.0, \mu_0^2 = 1.823405$$



$$\int_{-1}^1 dz \left( \frac{2}{1-z} \right)^{1/8} P_l(z)$$

$$\Rightarrow \frac{16}{7}, \frac{16}{35}, \frac{48}{161}, \frac{816}{3565}, \frac{12240}{64883}, \frac{493680}{3049501}, \frac{33456}{234577}, \frac{55760}{435643}, \frac{3602096}{30930653}, \frac{20129360}{187963199}, \frac{541373840}{5450932771}, \dots$$

$$\Delta_\sigma = \eta/2 = 1/8 \simeq 0.128$$




# EXACT FOUR POINT FUNCTION

$$g(u, v) = \frac{\langle \phi_1 \phi_2 \phi_3 \phi_4 \rangle}{\langle \phi_1 \phi_3 \rangle \langle \phi_2 \phi_4 \rangle}$$



$$= \frac{1}{\sqrt{2} |z|^{1/4} |1-z|^{1/4}} \left[ |1 + \sqrt{1-z}| + |1 - \sqrt{1-z}| \right]$$

$$u = z\bar{z} \quad , \quad v = (1-z)(1-\bar{z})$$

Crossing Sym:  $|1 + \sqrt{1-z}| + |1 - \sqrt{1-z}| = \sqrt{2 + 2\sqrt{(1-z)(1-\bar{z})} + 2\sqrt{z\bar{z}}}$



OPE Expansion:  $\phi \times \phi = \mathbf{1} + \phi^2$  or  $\sigma \times \sigma = \mathbf{1} -$



$$G_s(r, \theta) \propto 1 + \lambda_\epsilon^2 g_{\epsilon,0}(r, \theta) + \lambda_T^2 g_{T,2}(r, \theta)$$

$$\lambda_T^2 = \frac{\Delta_\sigma^2 d^2 |z|^{d-2}}{C_T (d-1)^2} \rightarrow \frac{1}{16C_T} \quad \text{for } d=2, \quad g_{T,2}(z) = -3 \left( 1 + \frac{1}{z} \left( 1 - \frac{z}{2} \right) \log(1-z) \right) + \text{c.c.}$$

# Fit TO OPE EXPANSION

$\mu^2$	$s$	$r_{\min} \leq r \leq r_{\max}$	norm	$\Delta_\epsilon$	$\lambda_\epsilon^2$	$c$
1.82241	9	$0.25 \leq r \leq 0.75$	0.2900	1.075	0.2536	0.4668
1.82241	9	$0.30 \leq r \leq 0.70$	0.2901	1.075	0.2533	0.4704
1.82241	9	$0.35 \leq r \leq 0.65$	0.2902	1.077	0.2533	0.4738
1.82241	9	$0.40 \leq r \leq 0.60$	0.2902	1.016	0.2427	0.4747
1.82241	18	$0.25 \leq r \leq 0.75$	0.2051	1.068	0.2563	0.4866
1.82241	18	$0.30 \leq r \leq 0.70$	0.2051	1.056	0.2544	0.4878
1.82241	18	$0.35 \leq r \leq 0.65$	0.2051	1.050	0.2535	0.4904
1.82241	18	$0.40 \leq r \leq 0.60$	0.2051	1.046	0.2526	0.4884
1.82241	36	$0.25 \leq r \leq 0.75$	0.1457	1.031	0.2528	0.4926
1.82241	36	$0.30 \leq r \leq 0.70$	0.1458	1.026	0.2519	0.4932
1.82241	36	$0.35 \leq r \leq 0.65$	0.1458	1.018	0.2508	0.4931
1.82241	36	$0.40 \leq r \leq 0.60$	0.1458	1.007	0.2486	0.4933

# CONCLUSION

1. *Simplicial Method for Scalars, Fermion and Gauge fields* **YES!**
2. *Quantum Corrections (QFE): Super renormalizable* **YES ?**
3. *4D UV asymptotical free: Local RG approach/Wilson Flow* **??**
4. **BSM** *Multi-flavor Gauge Theories with IR fixed point.* **WHY NOT**
5. *Simplicial Lattice in Anti de Sitter space* **YES! WHY?**
6. *General Mathematical “Proofs”* **DIFFICULT!**

# BACK UP SLIDES

# NEED COLLABORATORS & SUPPORT



# SPHERES AND CYLINDERS ARE NICE\*

- **Spheres** and **Cylinders** are Weyl trans\* & CFT are “preserved”.
- **Sphere:** For CFT, no finite volume approx & define: “c-theorems”
- **Cylinders:** Radial Quantized: Bndry of global AdS (H = Dilatations)



$$\mathbb{R}^d \rightarrow \mathbb{S}^d$$

$$ds_{flat}^2 = \sum_{\mu=1}^d dx^\mu dx^\mu = e^{\sigma(x)} d\Omega_d^2 \xrightarrow{Weyl} d\Omega_d^2.$$

$$\mathbb{R}^{d+1} \rightarrow \mathbb{R} \times \mathbb{S}^d$$

$$ds_{flat}^2 = \sum_{\mu=1}^{d+1} dx^\mu dx^\mu = e^{2t} (dt^2 + d\Omega_d^2) \xrightarrow{Weyl} (dt^2 + d\Omega_d^2).$$

$\mathbb{R}^d$

$\mathbb{S}^d$

$\mathbb{H}^d$


\*MAX SYM SPACE ARE NICE: Flat, de Sitter AdS

# ORIGINAL MOTIVATION: Radial Quantization

Conformal (near conformal) for

- BSM composite Higgs
- AdS/CFT weak-strong duality
- Model building & Critical Phenomena in general

$$\mathbb{R} \times \mathbb{T}^3 \quad \text{vs} \quad \mathbb{R} \times \mathbb{S}^3$$



Potential advantage: Scales increases exponentially in lattice size  $L$ !

$$1 < t < aL \implies 1 < \tau = \log(r) < L$$



# EXAMPLE SCALAR THEORY

$$S = \frac{1}{2} \int_{\mathcal{M}} d^D x \sqrt{g} [g^{\mu\nu} \partial_\mu \phi(x) \partial_\nu \phi(x) + m^2 \phi^2(x) + \lambda \phi^4(x)]$$



$$\vec{y} = \xi_1 \vec{r}_1 + \cdots + \xi_{D+1} \vec{r}_{D+1}$$

with  $\xi_1 + \cdots + \xi_{D+1} = 1$

$$\begin{aligned} I_\sigma &= \frac{1}{2} \int_\sigma d^D y [\vec{\nabla} \phi(y) \cdot \vec{\nabla} \phi(y) + m^2 \phi^2(y) + \lambda \phi^4(y)] \\ &= \frac{1}{2} \int_\sigma d^D \xi \sqrt{g} [g^{ij} \partial_i \phi(\xi) \partial_j \phi(\xi) + m^2 \phi^2(\xi) + \lambda \phi^4(\xi)] \end{aligned}$$



$$I_\sigma \simeq \sqrt{g_0} \left[ g_0^{ij} \frac{(\phi_i - \phi_0)(\phi_j - \phi_0)}{l_{i0} l_{j0}} + m^2 \phi_0^2 + \lambda \phi_0^4 \right]$$

# Using Binder Cumulants

In infinite volume

$U_{2n}=0$  in disordered phase

$U_{2n}=1$  in ordered phase

$0 < U_{2n} < 1$  on critical surface

$$U_4 = \frac{3}{2} \left( 1 - \frac{m_4}{3 m_2^2} \right) \quad m_n = \langle \phi^n \rangle$$

$$U_6 = \frac{15}{8} \left( 1 + \frac{m_6}{30 m_2^3} - \frac{m_4}{2 m_2^2} \right)$$

$$U_8 = \frac{315}{136} \left( 1 - \frac{m_8}{630 m_2^4} + \frac{2 m_6}{45 m_2^3} + \frac{m_4^2}{18 m_2^4} - \frac{2 m_4}{3 m_2^2} \right)$$

$$U_{10} = \frac{2835}{992} \left( 1 + \frac{m_{10}}{22680 m_2^5} - \frac{m_8}{504 m_2^4} - \frac{m_6 m_4}{108 m_2^5} + \frac{m_6}{18 m_2^3} + \frac{5 m_4^2}{36 m_2^4} - \frac{5 m_4}{6 m_2^2} \right)$$

$$U_{12} = \frac{155925}{44224} \left( 1 - \frac{m_{12}}{1247400 m_2^6} + \frac{m_{10}}{18900 m_2^5} + \frac{m_8 m_4}{2520 m_2^6} - \frac{m_8}{420 m_2^4} \right. \\ \left. + \frac{m_6^2}{2700 m_2^6} - \frac{m_6 m_4}{45 m_2^5} + \frac{m_6}{15 m_2^3} - \frac{m_4^3}{108 m_2^6} + \frac{m_4^2}{4 m_2^4} - \frac{m_4}{m_2^2} \right)$$

- $U_{2n,cr}$  are universal quantities.
- Deng and Blöte (2003):  $U_{4,cr}=0.851001$
- Higher critical cumulants computable using conformal  $2n$ -point functions:  
Luther and Peschel (1975)  
Dotsenko and Fateev (1984)

